

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

<b>1. REPORT DATE (DD-MM-YYYY)</b>		<b>2. REPORT TYPE</b> Technical Paper		<b>3. DATES COVERED (From - To)</b>	
<b>4. TITLE AND SUBTITLE</b>				<b>5a. CONTRACT NUMBER</b> F04611-99-C-0025	
				<b>5b. GRANT NUMBER</b>	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>				<b>5d. PROJECT NUMBER</b> 1011	
				<b>5e. TASK NUMBER</b> 001V	
				<b>5f. WORK UNIT NUMBER</b> 549882	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>				<b>8. PERFORMING ORGANIZATION REPORT</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>  Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b>  Approved for public release; distribution unlimited.					
<b>13. SUPPLEMENTARY NOTES</b>					
<b>14. ABSTRACT</b>  <div style="text-align: right; font-size: 2em; font-weight: bold;">20030128 109</div>					
<b>15. SUBJECT TERMS</b>					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>  A	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b> Leilani Richardson
<b>a. REPORT</b> Unclassified	<b>b. ABSTRACT</b> Unclassified	<b>c. THIS PAGE</b> Unclassified			<b>19b. TELEPHONE NUMBER</b> (include area code) (661) 275-5015

99-C0025  
549882  
1V  
MEMORANDUM FOR PRS (In-House Contractor Publication)

FROM: PROI (STINFO)

29 April 2002

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2002-088**  
Karl Christe et al., "Triphenylmethyldifluoramine: A Stable Reagent for the Synthesis of *gem*-  
Bis(difluoramines)"

J.C.S. Chemical Communications  
(Deadline: N/A)

(Statement A)

COMMUNICATION

ChemComm  
www.rsc.org/chemcomm

# Triphenylmethyldifluoramine: A Stable Reagent for the Synthesis of *gem*-Bis(difluoramines)

G. K. Surya Prakash,<sup>a\*</sup> Markus Etzkorn,<sup>a</sup> George A. Olah,<sup>a\*</sup> Karl O. Christe,<sup>a,b</sup> Stefan Schneider<sup>a</sup> and Ashwani Vij<sup>b</sup>


<sup>a</sup> Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, University Park, Los Angeles, CA 90089-1661, USA. Fax: +1 213 740 6270; Tel: +1 213 740 5984; E-mail: gprakash@usc.edu

<sup>b</sup> Air Force Research Laboratory, Edwards Air Force Base, CA 93524, USA.


This submission was created using the RSC ChemComm Template (DO NOT DELETE THIS TEXT)  
(LINE INCLUDED FOR SPACING ONLY - DO NOT DELETE THIS TEXT)

The conversion of ketones into geminal bis(difluoramines) can be achieved under mild two phase reaction conditions by employing triphenylmethyldifluoramine as an in situ source of difluoramine.

The introduction of a difluoramine functionality into organic moieties and the properties of the obtained target compounds have been investigated intensively.<sup>1</sup> More recently a new class of high-energy containing materials, *gem*-bis(difluoramine)-substituted heterocyclic nitramines has gained much attention as high-energy oxidizers: HNF<sub>x</sub> (1)<sup>2</sup> and TNFX (2)<sup>3</sup> have been successfully synthesized from their corresponding ketone derivatives. Usually this transformation is achieved under strongly acidic conditions with an excess of condensed difluoramine (HNF<sub>2</sub>)<sup>4</sup>, an unpredictably shock-sensitive and thermally unstable, gaseous compound.<sup>5</sup> HNF<sub>2</sub> can be generated from different precursors, e.g. tetrafluorohydrazine<sup>6</sup>, N,N-difluorourea<sup>7</sup> or N,N-difluorocarbamates.<sup>8</sup> A short report on the synthesis of triphenylmethyldifluoramine (3) mentions its hydrolysis to HNF<sub>2</sub>.<sup>9</sup> Except for 3, a stable crystalline compound, all of the other HNF<sub>2</sub> precursors have only limited stability.




Compound 3 proved to be a convenient reagent for the conversion of ketones (A) into *gem*-bis(difluoramines) (B) thus circumventing the potential hazards associated with previous methodologies.



Triphenylmethyldifluoramine (3) is accessible from chlorotriphenylmethane in multi-gram quantities as stable, non-hygroscopic, colorless crystals.<sup>9</sup> It can be stored without decomposition or any other hazards at ambient conditions. The complete characterization of 3 was carried out by IR, Raman, <sup>1</sup>H, <sup>13</sup>C and <sup>19</sup>F NMR spectroscopy.<sup>8</sup> Additionally, we obtained single crystals suitable for a *single crystal* X-ray crystal structure determination.<sup>8</sup>

The structural features of 3 are similar to those reported for *p*-bromophenyldiphenyl-methyldifluoramine.<sup>10</sup> The C-N distance of 1.523(2) Å found in 3 is longer than the corresponding distance

of 1.481 Å in triphenylmethyldifluoramine.<sup>11</sup> The presence of two highly electronegative fluorines on the nitrogen atom reduces back-donation from the nitrogen lone pair, thereby decreasing the C-N bond order. The torsion angles from the difluoramine group to the *ipso*-carbon atoms of the phenyl rings, namely F1-N1-C1-C8 and F2-N1-C1-C2 are 56.4(1)° and 39.1(1)°, respectively, and reflect the deviation of these atoms from perfect staggering.



The closest intramolecular H...F contacts found in the crystal structure of 3 result from H3...F1 and F2...F15 at 2.281(16) and 2.334(16) Å, respectively. The closest intermolecular H...N contacts result from H15...N1 and H9...N1 at 2.408(16) and 2.553(16) Å, respectively. All these contacts are shorter than the sum of the van der Waals radii of nitrogen and hydrogen (1.55 + 1.20 = 2.75 Å) or fluorine and hydrogen (1.47 + 1.20 = 2.67 Å). A plot of the unit cell is provided with the supplementary information.<sup>1,8</sup>

We have investigated two-phase organic solvent / oleum media bis(difluoraminations) with 3 under mild reaction conditions for some model ketones. Typically a solution of the ketone (A) and 2.2 equivalents of 3 in CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub> or CCl<sub>4</sub> were added dropwise to the same volume of oleum (30%) at 0 °C. Due to the potential hazards of the intermediate HNF<sub>2</sub> and any labile products, all reactions were carried out on a small scale (~1 mmol). The completion of the reactions was confirmed by NMR spectroscopy (<sup>19</sup>F, <sup>1</sup>H, <sup>13</sup>C). Thus, they were usually worked up after 15 to 20 min by pouring into ice/water, followed by extraction (CH<sub>2</sub>Cl<sub>2</sub>) and evaporation of the solvent from the dried organic phase.

**DISTRIBUTION STATEMENT A**  
Approved for Public Release  
Distribution Unlimited

Table 1


ENTRY	KETONE (A)	gem-BIS(DIFLUORAMINE) (B)	YIELD [%]	LITERATURE
1	Acetone (4)	2,2-Bis(difluoramino)propane (12)	90	4a
2	Cyclopentanone (5)	1,1-Bis(difluoramino)cyclopentane (13)	80	4a
3	Cyclohexanone (6)	1,1-Bis(difluoramino)cyclohexane (14)	80	4a
4	Cyclohexa-1,4-dione (7)	1,1,4,4-Tetrakis(difluoramino)cyclohexane (15)	85	4a
5	N-Acetyl-piperidinone (8)	4,4-Bis(difluoramino)-N-acetyl-piperidine (16)	85	14
6	9a (n = 1)	17a	ab	-
7	9b (n = 9)	17b	80 <sup>b</sup>	-
8	9c (n = 10)	17c	85 <sup>b</sup>	-
9	10	18	90 <sup>b</sup>	-
10	11a (R = H)	20a	0 <sup>c</sup>	-
11	11b (R = CH <sub>3</sub> )	20b	0 <sup>c</sup>	-

<sup>a</sup> decomposition upon work-up; <sup>b</sup> analytical data provided in the supplementary information<sup>§</sup>; <sup>c</sup> only formation of the cage-systems 19a (19b)<sup>§</sup> in ca. 80% yield was observed

The crude product contained only triphenylmethanol and the gem-bis(difluoramino) that (on a larger scale) can be purified by repeated crystallization, chromatography on neutral aluminum oxide or distillation, respectively. The given yields are based on <sup>1</sup>H NMR data of the crude product containing only the gem-bis(difluoramino) (B) and triphenylmethanol. A variety of known (entry 1-5) as well as new (entry 6-11) difluoramino derivatives was prepared in good to excellent yields.

neat HNF<sub>2</sub> and therefore minimizes its potential hazards. Work is underway to further investigate the reaction mechanism, find alternative preparations of the reagent, and synthesize potential high-energy materials.

We gratefully acknowledge the financial support by the Office of Naval Research (Program Manager: Dr. Judah Goldwasser), the Air Force Office of Scientific Research and the National Science Foundation.



Although 17a could not yet be isolated on this small scale, it represents the smallest alicyclic geminal bis(difluoramino) so far characterized. The bicyclic 11a (11b) could only be converted to the bridged hemiaminal 19a (19b); neither a larger excess of 3 nor extended reaction time did furnish the tetrakis(difluoramino) 20a (20b).

Mechanistically, the reactions involve cleavage of 3 to the triphenylmethyl cation and difluoramino. The latter or the in situ product, difluorsulfamic acid reacts with the ketone A to provide gem-bis(difluoramino) B. It is noteworthy that our reaction did not proceed in conc. H<sub>2</sub>SO<sub>4</sub>, although acidolysis of 3 was observed spectroscopically: the <sup>19</sup>F NMR signal of 3 disappeared and the trityl cation was observed in the <sup>13</sup>C NMR spectrum. Also employing other strong acids such as FSO<sub>3</sub>H, CF<sub>3</sub>CO<sub>2</sub>H, CF<sub>3</sub>SO<sub>3</sub>H or CF<sub>3</sub>SO<sub>3</sub>H/(CF<sub>3</sub>SO<sub>2</sub>)<sub>2</sub>O resulted in similar behavior. In the case of FSO<sub>3</sub>H/SbF<sub>5</sub>, only decomposition of 3 was observed. We have never been able to observe free or protonated difluoramino.<sup>12</sup> It is also possible that under the reaction conditions SO<sub>3</sub> in oleum – besides being a water scavenger – might react with 3 to form F<sub>2</sub>NSO<sub>3</sub>H as the ultimate difluoramino reagent (vide supra). The triphenylmethyl cation might also act like a Lewis acid catalyst, polarizing the carbonyl group (Mukaiyama type activation<sup>13</sup>).

In conclusion, the readily preparable, stable triphenylmethyldifluoramino (3) can be conveniently employed for the efficient conversion of ketones (A) under mild reaction conditions to the respective geminal bis(difluoramines) (B). This is the first difluoramination methodology that avoids the use of

## Notes and references

§ Supplementary information contains analytical data for 3, 18, 19, 20, 21 (2 pages) and a plot of the unit cell of 3 (3 pages).

† Crystal data: 3, C<sub>10</sub>H<sub>12</sub>F<sub>2</sub>N, M<sub>r</sub> = 295.32, monoclinic, a = 10.3882(12), b = 9.8259(11), c = 14.6065(16) Å, β = 92.402(2)°, V = 1489.6(3) Å<sup>3</sup>, T = 213(2) K, space group P2(1)/n, Z = 4, D<sub>c</sub> = 1.317 Mg m<sup>-3</sup>, μ (Mo-Kα) = 0.094 mm<sup>-1</sup>, 9649 reflections measured, 3035 unique (R<sub>int</sub> = 0.0321). The final wR2 was 0.1098 (all data).

- W. Hanefeld, *N,N*-Dihalogen-amine, ed. D. Klamann in *Methoden der Organischen Chemie (Houben-Weyl) Nitrogen Compounds*, Vol. E16a / Part 2, Georg-Thieme Verlag, Stuttgart, New York, 1990, pp. 893-925 and references therein.
- (a) R. D. Chapman, M. F. Welker, C. B. Kreutzberger, *J. Org. Chem.*, 1998, 63, 1566; (b) R. D. Chapman, R. D. Gilardi, M. F. Welker, C. B. Kreutzberger, *J. Org. Chem.*, 1999, 64, 960.
- T. Axenrod, X.-P. Guan, J. Sun, L. Qi, R. D. Chapman, R. D. Gilardi, *Tetrahedron Lett.*, 2001, 42, 2621.
- (a) K. Baum, *J. Amer. Chem. Soc.*, 1968, 90, 7083; (b) W. H. Graham, J. P. Freeman, *J. Org. Chem.*, 1969, 34, 2589; (c) A. V. Fokin, Y. M. Kosyrev, V. A. Makarov, N. P. Novoselov, *Dokl. Akad. Nauk SSSR*, 1969, 186, 112.
- E. W. Lawless, I. C. Smith, *Inorganic High Energy Oxidizers (Synthesis, Structure and Properties)*, Marcel Dekker Inc., New York, NY, USA, 1968, 69.
- J. P. Freeman, A. Kennedy, C. B. Colburn, *J. Amer. Chem. Soc.*, 1960, 82, 5304.
- (a) V. Grakauskas, K. Baum, *J. Amer. Chem. Soc.*, 1970, 92, 2096; (b) R. E. Banks, R. N. Hazeldine, J. P. Lallu, *J. Chem. Soc. (c)*, 1966, 1514.
- (a) V. Grakauskas, K. Baum, *J. Amer. Chem. Soc.*, 1969, 91, 1679; (b) J. M. J. Lambert, G. M. Alain (Societe Nationale des Poudres et Explosives), 1973, DE 2320119; (c) D. L. Klapotek, G. B. Hobrock, *Inorg. Chem.*, 1967, 6, 1751.
- (a) R. C. Petry, J. P. Freeman, *J. Amer. Chem. Soc.*, 1961, 83, 3912; (b) W. H. Graham, C. O. Parker, *J. Org. Chem.*, 1963, 28, 850.
- J. R. Surlis, C. I. Bumgardner, J. Bordner, *J. Fluorine Chem.*, 1975, 5, 467.
- C. Gidwell, G. Furgeson, *Acta Crystallogr., Sec. C Crst. Str. Commun.*, 1994, 50, 924.
- K. O. Christie, *Inorg. Chem.*, 1975, 14, 2821.
- S. Kobayashi, M. Murakami, T. Mukaiyama, *Chem. Lett.*, 1985, 1535.
- J. Zhang, J. Oxley, J. Smith, C. Bedford, R. Chapman, *J. Mass Spectrometry*, 2000, 35, 841.